arXiv:1503.04893v1 [math.NT] 17 Mar 2015

SOME IDENTITIES OF q-BERNOULLI POLYNOMIALS UNDER SYMMETRY GROUP S_3

DMITRY V. DOLGY, DAE SAN KIM, AND TAEKYUN KIM

Abstract. In this paper, we give some new identities of Carlitz q-Bernoulli polynomials under symmetry group S_3. The derivatives of identities are based on the q-Volkenborn integral expression of the generating function for the Carlitz q-Bernoulli polynomials and the q-Volkenborn integral equations that can be expressed as the exponential generating functions for the q-power sums.

1. Introduction

Let p be a fixed prime number. Throughout this paper, $\mathbb{Z}_p$, $\mathbb{Q}_p$ and $\mathbb{C}_p$ will, respectively, denote the ring of p-adic integers, the field of p-adic rational numbers and the completion of algebraic closure of $\mathbb{Q}_p$. Let ν_p be the normalized exponential valuation of $\mathbb{C}_p$ with $|p|_p = p^{-\nu_p(p)} = \frac{1}{p}$ and let q be an indeterminate in $\mathbb{C}_p$ with $|1-q|_p < p^{-\frac{1}{p-1}}$. The q-number of x is defined as $[x]_q = \frac{1-q^x}{1-q}$. Let $UD(\mathbb{Z}_p)$ be the space of uniformly differentiable functions on $\mathbb{Z}_p$. For $f \in UD(\mathbb{Z}_p)$, the q-Volkenborn integral on $\mathbb{Z}_p$ is defined by Kim to be

$$I_q(f) = \int_{\mathbb{Z}_p} f(x)\, d\mu_q(x) = \lim_{N\to\infty} \frac{1}{[p^N]_q} \sum_{x=0}^{p^N-1} f(x)\, q^x, \quad (\text{see [13]}). \tag{1.1}$$

In (1.1), we note that

$$qI_q(f_1) - I_q(f) = \frac{q-1}{\log q} f'(0) + (q-1) f(0), \tag{1.2}$$

where $f_1(x) = f(x+1)$.

In general, one derives

$$q^n I_q(f_n) - I_q(f) = \frac{q-1}{\log q} \sum_{l=0}^{n-1} f'(l)\, q^l + (q-1) \sum_{l=0}^{n-1} f(l)\, q^l, \tag{1.3}$$

where $f_n(x) = f(x+n)$, $(n \geq 0)$, (see [15, 16, 27]).

It is well known that the Bernoulli numbers are given by

$$B_0 = 1, \quad (B+1)^n - B_n = \delta_{1,n}, \quad (\text{see [1–27]}) \tag{1.4}$$

with the usual convention about replacing B^n by B_n.

The Bernoulli polynomials are defined by

$$B_n(x) = \sum_{l=0}^{n} \binom{n}{l} B_l x^{n-l}, \quad (n \geq 0), \quad (\text{see [29, 30]}). \tag{1.5}$$

2000 *Mathematics Subject Classification.* 05A19; 11B65; 11B68.

Key words and phrases. Carlitz q-Bernoulli polynomial; q-Volkenborn integral.

In [4, 5], L. Carlitz considered a q-analogue of Bernoulli numbers as follows :

$$\beta_{0,q}=1, \quad q\left(q\beta_q+1\right)^n-\beta_{n,q}=\begin{cases}1 & \text{if } n=1,\\ 0 & \text{if } n>1,\end{cases} \tag{1.6}$$

with the usual convention about replacing β_q^i by $\beta_{i,q}$.

He also defined q-Bernoulli polynomials as follows :

$$\beta_{n,q}\left(x\right)=\sum_{l=0}^{n}\binom{n}{l}q^{lx}\beta_{l,q}\left[x\right]_q^{n-l}, \quad \left(\text{see [4, 5]}\right). \tag{1.7}$$

From (1.6), Carlitz derived the following equation :

$$\beta_{n,q}=\frac{1}{\left(1-q\right)^n}\sum_{l=0}^{n}\binom{n}{l}\left(-1\right)^l\frac{l+1}{\left[l+1\right]_q}, \tag{1.8}$$

and

$$\beta_{n,q}\left(x\right)=\frac{1}{\left(1-q\right)^n}\sum_{l=0}^{n}\binom{n}{l}\left(-1\right)^l q^{lx}\frac{l+1}{\left[l+1\right]_q}, \quad \left(\text{see [4]}\right). \tag{1.9}$$

Carlitz q-Bernoulli numbers and polynomials are also given by q-Volkenborn integrals on $\mathbb{Z}_p$ due to T. Kim [13] :

$$\begin{aligned}\sum_{n=0}^{\infty}\beta_{n,q}\frac{t^n}{n!} &= \int_{\mathbb{Z}_p}e^{[y]_q t}d\mu_q\left(y\right)\\ &= \sum_{m=0}^{\infty}q^m e^{[m]_q t}\left(1-q-q^m t\right),\end{aligned} \tag{1.10}$$

and

$$\begin{aligned}\sum_{n=0}^{\infty}\beta_{n,q}\left(x\right)\frac{t^n}{n!} &= \int_{\mathbb{Z}_p}e^{[x+y]_q t}d\mu_q\left(y\right)\\ &= \sum_{m=0}^{\infty}q^m e^{[x+m]_q t}\left(1-q-q^{x+m}t\right).\end{aligned} \tag{1.11}$$

The purpose of this paper is to give some new identities of Carlitz q-Bernoulli polynomials under symmetry group S_3. The derivations of identities are based on the q-Volkenborn integral expression of the generating function for the Carlitz q-Bernoulli polynomials and the q-Volkenborn integrals equations that can be expressed as the exponential generating functions for the q-power sums.

2. Some identities of Carlitz q-Bernoulli polynomials

By (1.7), we easily get

$$\begin{aligned}\beta_{n,q}\left(x+y\right) &= \sum_{l=0}^{n}\binom{n}{l}q^{lx}\beta_{l,q}\left(y\right)\left[x\right]_q^{n-l}\\ &= \sum_{l=0}^{n}\binom{n}{l}q^{(n-l)x}\beta_{n-l,q}\left(y\right)\left[x\right]_q^{l}.\end{aligned} \tag{2.1}$$

On the other hand, Carlitz also introduced the expression of q-Bernoulli polynomials $\beta_{n,q}^{(h,k)}(x)$ as follows :

$$\beta_{n,q}^{(h,k)}(x)=\frac{1}{(1-q)^n}\sum_{j=0}^{n}\binom{n}{j}(-1)^j q^{jx}\frac{(j+h)_k}{[j+h]_{q,k}}, \tag{2.2}$$

where

$$[j+h]_{q,k}=[j+h]_q[j+h-1]_q\cdots[j+h-k+1]_q, \tag{2.3}$$

and

$$(j+h)_k=(j+h)(j+h-1)\cdots(j+h-k+1),\quad \text{(see [5, 13])}. \tag{2.4}$$

T. Kim [13] obtained the Witt-type formula for $\beta_{n,q}^{(h,k)}(x)$ as follows :

$$\beta_{n,q}^{(h,k)}(x)=\int_{\mathbb{Z}_p}\cdots\int_{\mathbb{Z}_p}q^{\sum_{l=1}^{k}(h-l)y_l}[x+y_1+\cdots+y_k]_q^n\,d\mu_q(y_1)\cdots d\mu_q(y_k). \tag{2.5}$$

If $k=1$, then $\beta_{n,q}^{(h,1)}(x)$ will be simply denoted by $\beta_{n,q}^{(h)}(x)$ so that

$$\beta_{n,q}^{(h)}(x)=\int_{\mathbb{Z}_p}q^{(h-1)y}[x+y]_q^n\,d\mu_q(y). \tag{2.6}$$

The following simple facts will be used over and over again :

$$[a+b]_q=[a]_q+q^a[b]_q. \tag{2.7}$$

By (2.7), we easily see that

$$[a+b+c]_q=[a]_q+q^a[b]_q+q^{a+b}[c]_q, \tag{2.8}$$

and

$$[ab]_q=[a]_q[b]_{q^a}. \tag{2.9}$$

First, we will consider the following triple integral which is obviously invariant under any permutations of w_1, w_2, w_3.

So the expression obtained from this after integration will also be invariant under any permutations of w_1, w_2, w_3. This observation is simple enough but it is the philosophy that underlies this paper.

$$\begin{aligned}I=\int_{\mathbb{Z}_p^3}&e^{[w_2w_3x_1+w_1w_3x_2+w_1w_2x_3+w_1w_2w_3(y_1+y_2+y_3)]_q t}\\&\times d\mu_{q^{w_2w_3}}(x_1)\,d\mu_{q^{w_1w_3}}(x_2)\,d\mu_{q^{w_1w_2}}(x_3).\end{aligned} \tag{2.10}$$

It is easy to show that

$$\begin{aligned}&[w_2w_3x_1+w_1w_3x_2+w_1w_2x_3+w_1w_2w_3(y_1+y_2+y_3)]_q\\&=[w_2w_3]_q[x_1+w_1y_1]_{q^{w_2w_3}}+q^{w_2w_3(x_1+w_1y_1)}[w_1w_3]_q[x_2+w_2y_2]_{q^{w_1w_3}}\\&\quad+q^{w_2w_3(x_1+w_1y_1)+w_1w_3(x_2+w_2y_2)}[w_1w_2]_q[x_3+w_3y_3]_{q^{w_1w_2}}.\end{aligned} \tag{2.11}$$

So the integrand is

$$
\begin{aligned}
& e^{[w_2w_3x_1+w_1w_3x_2+w_1w_2x_3+w_1w_2w_3(y_1+y_2+y_3)]_q t} \\
=& e^{[w_2w_3]_q [x_1+w_1y_1]_{q^{w_2w_3}} t} e^{q^{w_2w_3(x_1+w_1y_1)}[w_1w_3]_q [x_2+w_2y_2]_{q^{w_1w_3}} t} \\
& \times e^{q^{w_2w_3(x_1+w_1y_1)+w_1w_3(x_2+w_2y_2)}[w_1w_2]_q [x_3+w_3y_3]_{q^{w_1w_2}} t} \\
=& \sum_{n=0}^{\infty} \sum_{k+l+m=n} \binom{n}{k,l,m} [w_2w_3]_q^k [w_1w_3]_q^l [w_1w_2]_q^m q^{w_1w_2w_3(l+m)y_1} \\
& \times q^{w_1w_2w_3my_2} q^{w_2w_3(l+m)x_1} [x_1+w_1y_1]_{q^{w_2w_3}}^k q^{w_1w_3mx_2} \\
& \times [x_2+w_2y_2]_{q^{w_1w_3}}^l [x_3+w_3y_3]_{q^{w_1w_2}}^m \frac{t^n}{n!}.
\end{aligned}
\tag{2.12}
$$

Thus the integral in (2.10) is given by

$$
\begin{aligned}
I =& \sum_{n=0}^{\infty} \left\{ \sum_{k+l+m=n} \binom{n}{k,l,m} [w_2w_3]_q^k [w_1w_3]_q^l [w_1w_2]_q^m q^{w_1w_2w_3(l+m)y_1} \right. \\
& \times q^{w_1w_2w_3my_2} \int_{\mathbb{Z}_p} q^{w_2w_3(l+m)x_1} [x_1+w_1y_1]_{q^{w_2w_3}}^k d\mu_{q^{w_2w_3}}(x_1) \\
& \times \int_{\mathbb{Z}_p} q^{w_1w_3mx_2} [x_2+w_2y_2]_{q^{w_1w_3}}^l d\mu_{q^{w_1w_3}}(x_2) \\
& \left. \times \int_{\mathbb{Z}_p} [x_3+w_3y_3]_{q^{w_1w_2}}^m d\mu_{q^{w_1w_2}}(x_3) \right\} \frac{t^n}{n!} \\
=& \sum_{n=0}^{\infty} \left\{ \sum_{k+l+m=n} \binom{n}{k,l,m} [w_2w_3]_q^k [w_1w_3]_q^l \right. \\
& \times [w_1w_2]_q^m q^{w_1w_2w_3(l+m)y_1} q^{w_1w_2w_3my_2} \\
& \left. \times \beta_{k,q^{w_2w_3}}^{(l+m+1)}(w_1y_1) \beta_{l,q^{w_1w_3}}^{(m+1)}(w_2y_2) \beta_{m,q^{w_1w_2}}(w_3y_3) \right\} \frac{t^n}{n!}.
\end{aligned}
\tag{2.13}
$$

Thus, by (2.13), we get the following theorem.

Theorem 1. *Let w_1, w_2, w_3 be any positive integers, n any nonnegative integer. Then the following expression is invariant under any permutation of w_1, w_2, w_3 so that it gives us six symmetries :*

$$\begin{aligned}
&\sum_{k+l+m=n} \binom{n}{k,l,m} [w_2w_3]_q^k [w_1w_3]_q^l [w_1w_2]_q^m q^{w_1w_2w_3(l+m)y_1} \\
&\quad \times q^{w_1w_2w_3my_2} \beta_{k,q^{w_2w_3}}^{(l+m+1)}(w_1y_1) \beta_{l,q^{w_1w_3}}^{(m+1)}(w_2y_2) \beta_{m,q^{w_1w_2}}(w_3y_3) \\
=& \sum_{k+l+m=n} \binom{n}{k,l,m} [w_1w_3]_q^k [w_2w_3]_q^l [w_1w_2]_q^m q^{w_1w_2w_3(l+m)y_1} \\
&\quad \times q^{w_1w_2w_3my_2} \beta_{k,q^{w_1w_3}}^{(l+m+1)}(w_2y_1) \beta_{l,q^{w_2w_3}}^{(m+1)}(w_1y_2) \beta_{m,q^{w_1w_2}}(w_3y_3) \\
=& \sum_{k+l+m=n} \binom{n}{k,l,m} [w_1w_3]_q^k [w_1w_2]_q^l [w_2w_3]_q^m q^{w_1w_2w_3(l+m)y_1} \\
&\quad \times q^{w_1w_2w_3my_2} \beta_{k,q^{w_1w_3}}^{(l+m+1)}(w_2y_1) \beta_{l,q^{w_1w_2}}^{(m+1)}(w_3y_2) \beta_{m,q^{w_2w_3}}(w_1y_3) \\
=& \sum_{k+l+m=n} \binom{n}{k,l,m} [w_2w_3]_q^k [w_1w_2]_q^l [w_1w_3]_q^m q^{w_1w_2w_3(l+m)y_1} \\
&\quad \times q^{w_1w_2w_3my_2} \beta_{k,q^{w_2w_3}}^{(l+m+1)}(w_1y_1) \beta_{l,q^{w_1w_2}}^{(m+1)}(w_3y_2) \beta_{m,q^{w_1w_3}}(w_2y_3) \\
=& \sum_{k+l+m=n} \binom{n}{k,l,m} [w_1w_2]_q^k [w_2w_3]_q^l [w_1w_3]_q^m q^{w_1w_2w_3(l+m)y_1} \\
&\quad \times q^{w_1w_2w_3my_2} \beta_{k,q^{w_1w_2}}^{(l+m+1)}(w_3y_1) \beta_{l,q^{w_2w_3}}^{(m+1)}(w_1y_2) \beta_{m,q^{w_1w_3}}(w_2y_3) \\
=& \sum_{k+l+m=n} \binom{n}{k,l,m} [w_1w_2]_q^k [w_1w_3]_q^l [w_2w_3]_q^m q^{w_1w_2w_3(l+m)y_1} \\
&\quad \times q^{w_1w_2w_3my_2} \beta_{k,q^{w_1w_2}}^{(l+m+1)}(w_3y_1) \beta_{l,q^{w_1w_3}}^{(m+1)}(w_2y_2) \beta_{m,q^{w_2w_3}}(w_1y_3).
\end{aligned}$$

We define, for nonnegative integers $n, m, w, T_{n,m}(w|q)$ as

$$T_{n,m}(w|q) = \sum_{i=0}^{w} q^{ni} [i]_q^m . \tag{2.14}$$

In particular, for $w = 0$, we have

$$T_{n,m}(0|q) = \begin{cases} 1, & \text{if } m = 0 \\ 0, & \text{if } m > 0, \end{cases} \tag{2.15}$$

and

$$T_{n,0}(w|q) = \begin{cases} w+1 & \text{if } n = 0 \\ [w+1]_{q^n} & \text{if } n > 0. \end{cases} \tag{2.16}$$

From (1.3), we have

$$
\begin{aligned}
&\left(q^{w_1w_2}\right)^{w_3}\int_{\mathbb{Z}_p} e^{[w_1w_2(x+w_3)]_q t} d\mu_{q^{w_1w_2}}(x) \\
&-\int_{\mathbb{Z}_p} e^{[w_1w_2x]_q t} d\mu_{q^{w_1w_2}}(x) \\
=&t\left[w_1w_2\right]_q \sum_{i=0}^{w_3-1} q^{2w_1w_2i} e^{[w_1w_2i]_q t} \\
&+(q-1)\left[w_1w_2\right]_q \sum_{i=0}^{w_3-1} q^{w_1w_2i} e^{[w_1w_2i]_q t} \\
=&\sum_{m=0}^{\infty} T_{2,m}\left(w_3-1|q^{w_1w_2}\right)\left[w_1w_2\right]_q^{m+1}\frac{t^{m+1}}{m!} \\
&+(q-1)\sum_{m=0}^{\infty} T_{1,m}\left(w_3-1|q^{w_1w_2}\right)\left[w_1w_2\right]_q^{m+1}\frac{t^{m}}{m!}.
\end{aligned}
\tag{2.17}
$$

Thus, we have the following lemma.

Lemma 2. *For $w_1, w_2, w_3 \geq 1$, we have*

$$
\begin{aligned}
&q^{w_1w_2w_3}\int_{\mathbb{Z}_p} e^{[w_1w_2(x+w_3)]_q t} d\mu_{q^{w_1w_2}}(x) - \int_{\mathbb{Z}_p} e^{[w_1w_2x]_q t} d\mu_{q^{w_1w_2}}(x) \\
=&\sum_{m=0}^{\infty} \frac{\left[w_1w_2\right]_q^m}{m!} t^m \int_{\mathbb{Z}_p}\left(q^{w_1w_2w_3}\left[x+w_3\right]_{q^{w_1w_2}}^m-[x]_{q^{w_1w_2}}^m\right) d\mu_{q^{w_1w_2}}(x) \\
=&\sum_{m=0}^{\infty} T_{2,m}\left(w_3-1|q^{w_1w_2}\right)\left[w_1w_2\right]_q^{m+1}\frac{t^{m+1}}{m!} \\
&+(q-1)\sum_{m=0}^{\infty} T_{1,m}\left(w_3-1|q^{w_1w_2}\right)\left[w_1w_2\right]_q^{m+1}\frac{t^{m}}{m!} \\
=&t\left[w_1w_2\right]_q \sum_{i=0}^{w_3-1} q^{2w_1w_2i} e^{[w_1w_2i]_q t}+(q-1)\left[w_1w_2\right]_q \sum_{i=0}^{w_3-1} q^{w_1w_2i} e^{[w_1w_2i]_q t}.
\end{aligned}
$$

Now, we consider the following difference of triple integrals.

$$
\begin{aligned}
I_1 =&q^{w_1w_2w_3}\int_{\mathbb{Z}_p^3} e^{[w_2w_3x_1+w_1w_3x_2+w_1w_2x_3+w_1w_2w_3(y_1+y_2+1)]_q t} \\
&\times d\mu_{q^{w_2w_3}}(x_1)\, d\mu_{q^{w_1w_3}}(x_2)\, d\mu_{q^{w_1w_2}}(x_3) \\
&-\int_{\mathbb{Z}_p^3} e^{[w_2w_3x_1+w_1w_3x_2+w_1w_2x_3+w_1w_2w_3(y_1+y_2)]_q t} \\
&\times d\mu_{q^{w_2w_3}}(x_1)\, d\mu_{q^{w_1w_3}}(x_2)\, d\mu_{q^{w_1w_2}}(x_3),
\end{aligned}
\tag{2.18}
$$

which is obviously invariant under any permutations of w_1, w_2, w_3.

We put

$$
a=a\left(x_1\right)=q^{w_2w_3(x_1+w_1y_1)}, \quad b=b\left(x_2\right)=q^{w_1w_3(x_2+w_2y_2)}.
\tag{2.19}
$$

Then, by (2.18), we get

$$\begin{aligned} I_1 = &\sum_{k,l=0}^{\infty} [w_2w_3]_q^k [w_1w_3]_q^l \frac{t^{k+l}}{k!l!} \int_{\mathbb{Z}_p^2} a^l [x_1+w_1y_1]_{q^{w_2w_3}}^k [x_2+w_2y_2]_{q^{w_1w_3}}^l \\ &\times \left\{ \sum_{m=0}^{\infty} \frac{[w_1w_2]_q^m (abt)^m}{m!} \right. \\ &\times \left. \int_{\mathbb{Z}_p} \left(q^{w_1w_2w_3} [x_3+w_3]_{q^{w_1w_2}}^m - [x_3]_{q^{w_1w_2}}^m \right) d\mu_{q^{w_1w_2}}(x_3) \right\} \\ &\times d\mu_{q^{w_2w_3}}(x_1)\, d\mu_{q^{w_1w_3}}(x_2). \end{aligned} \tag{2.20}$$

From Lemma 2, the inner sum is

$$\begin{aligned} &\sum_{m=0}^{\infty} \frac{[w_1w_2]_q^{m+1} (abt)^{m+1}}{m!} T_{2,m}(w_3-1|q^{w_1w_2}) \\ &+ (q-1) \sum_{m=0}^{\infty} \frac{[w_1w_2]_q^{m+1} (abt)^m}{m!} T_{1,m}(w_3-1|q^{w_1w_2}). \end{aligned} \tag{2.21}$$

Thus, by (2.20) and (2.21), we get

$$\begin{aligned} I_1 = &\sum_{k,l,m=0}^{\infty} [w_2w_3]_q^k [w_1w_3]_q^l [w_1w_2]_q^{m+1} \frac{t^{k+l+m+1}}{k!l!m!} T_{2,m}(w_3-1|q^{w_1w_2}) \\ &\times \int_{\mathbb{Z}_p} a^{l+m+1} [x_1+m_1y_1]_{q^{w_2w_3}}^k d\mu_{q^{w_2w_3}}(x_1) \\ &\times \int_{\mathbb{Z}_p} b^{m+1} [x_2+w_2y_2]_{q^{w_1w_3}}^l d\mu_{q^{w_1w_3}}(x_2) \\ &+ (q-1) \sum_{k,l,m=0}^{\infty} [w_2w_3]_q^k [w_1w_3]_q^l [w_1w_2]_q^{m+1} \frac{t^{k+l+m}}{k!l!m!} \\ &\times T_{1,m}(w_3-1|q^{w_1w_2}) \int_{\mathbb{Z}_p} a^{l+m} [x_1+w_1y_1]_{q^{w_2w_3}}^k d\mu_{q^{w_2w_3}}(x_1) \\ &\times \int_{\mathbb{Z}_p} b^m [x_2+w_2y_2]_{q^{w_1w_3}}^l d\mu_{q^{w_1w_3}}(x_2). \end{aligned} \tag{2.22}$$

Recovering $a = q^{w_2w_3(x_1+w_1y_1)}$ and $b = q^{w_1w_3(x_2+w_2y_2)}$, (2.18) can be rewritten as :

$$\begin{aligned} I_1 = &\sum_{n=0}^{\infty} \left\{ \sum_{k+l+m=n} \binom{n}{k,l,m} [w_2w_3]_q^k [w_1w_3]_q^l [w_1w_2]_q^{m+1} \right. \\ &\times T_{2,m}(w_3-1|q^{w_1w_2})\, q^{w_1w_2w_3(l+m+1)y_1} q^{w_1w_2w_3(m+1)y_2} \\ &\times \int_{\mathbb{Z}_p} q^{w_2w_3(l+m+1)x_1} [x_1+w_1y_1]_{q^{w_2w_3}}^k d\mu_{q^{w_2w_3}}(x_1) \end{aligned} \tag{2.23}$$

$$\begin{aligned}
&\times \int_{\mathbb{Z}_p} q^{w_1w_3(m+1)x_2} \left[x_2+w_2y_2\right]_{q^{w_1w_3}}^{l} d\mu_{q^{w_1w_3}}\left(x_2\right) \Bigg\} \frac{t^{n+1}}{n!} \\
&+ (q-1)\sum_{n=0}^{\infty} \Bigg\{ \sum_{k+l+m=n} \binom{n}{k,l,m} \\
&\times \left[w_2w_3\right]_q^{k} \left[w_1w_3\right]_q^{l} \left[w_1w_2\right]_q^{m+1} T_{1,m}\left(w_3-1|q^{w_1w_2}\right)
\end{aligned}$$

$$\begin{aligned}
&\times q^{w_1w_2w_3(l+m)y_1} q^{w_1w_2w_3my_2} \\
&\times \int_{\mathbb{Z}_p} q^{w_2w_3(l+m)x_1} \left[x_1+w_1y_1\right]_{q^{w_2w_3}}^{k} d\mu_{q^{w_2w_3}}\left(x_1\right) \\
&\times \int_{\mathbb{Z}_p} q^{w_1w_3mx_2} \left[x_2+w_2y_2\right]_{q^{w_1w_3}}^{l} d\mu_{q^{w_1w_3}}\left(x_2\right) \Bigg\} \frac{t^{n}}{n!} \\
=&\sum_{n=0}^{\infty} \Bigg\{ \sum_{k+l+m=n} \binom{n}{k,l,m} \beta_{k,q^{w_2w_3}}^{(l+m+2)}\left(w_1y_1\right) \\
&\times \beta_{l,q^{w_1w_3}}^{(m+2)}\left(w_2y_2\right) T_{2,m}\left(w_3-1|q^{w_1w_2}\right) q^{w_1w_2w_3(l+m+1)y_1} \\
&\times q^{w_1w_2w_3(m+1)y_2} \left[w_2w_3\right]_q^{k} \left[w_1w_3\right]_q^{l} \left[w_1w_2\right]_q^{m+1} \Bigg\} \frac{t^{n+1}}{n!} \\
&+ (q-1)\sum_{n=0}^{\infty} \Bigg\{ \sum_{k+l+m=n} \binom{n}{k,l,m} \beta_{k,q^{w_2w_3}}^{(l+m+1)}\left(w_1y_1\right) \\
&\times \beta_{l,q^{w_1w_3}}^{(m+1)}\left(w_2y_2\right) T_{1,m}\left(w_3-1|q^{w_1w_2}\right) \\
&\times q^{w_1w_2w_3(l+m)y_1} q^{w_1w_2w_3my_2} \left[w_2w_3\right]_q^{k} \left[w_1w_3\right]_q^{l} \left[w_1w_2\right]_q^{m+1} \Bigg\} \frac{t^{n}}{n!}.
\end{aligned}$$

Separating the term corresponding to $n=0$ from the second term and after rearranging, we get :

$$\begin{aligned}
I_1 =& (q-1)\left[w_1w_2w_3\right]_q + \sum_{n=1}^{\infty} \Bigg\{ \sum_{k+l+m=n-1} \binom{n}{k,l,m} \\
&\times \beta_{k,q^{w_2w_3}}^{(l+m+2)}\left(w_1y_1\right) \beta_{l,q^{w_1w_3}}^{(m+2)}\left(w_2y_2\right) T_{2,m}\left(w_3-1|q^{w_1w_2}\right) \\
&\times q^{w_1w_2w_3(l+m+1)y_1} q^{w_1w_2w_3(m+1)y_2} \left[w_2w_3\right]_q^{k} \left[w_1w_3\right]_q^{l} \left[w_1w_2\right]_q^{m+1} \\
&\quad + (q-1) \sum_{k+l+m=n} \binom{n}{k,l,m} \beta_{k,q^{w_2w_3}}^{(l+m+1)}\left(w_1y_1\right) \\
&\quad \times \beta_{l,q^{w_1w_3}}^{(m+1)}\left(w_2y_2\right) T_{1,m}\left(w_3-1|q^{w_1w_2}\right) \\
&\quad \times q^{w_1w_2w_3(l+m)y_1} q^{w_1w_2w_3my_2} \\
&\quad \times \left[w_2w_3\right]_q^{k} \left[w_1w_3\right]_q^{l} \left[w_1w_2\right]_q^{m+1} \Bigg\} \frac{t^{n}}{n!}.
\end{aligned} \tag{2.24}$$

As this expression is invariant under any permutations in w_1, w_2, w_3, we get the following theorem.

Theorem 3. *Let $w_1, w_2, w_3 \in \mathbb{Z}$ with $w_1 \geq 1$, $w_2 \geq 1$, $w_3 \geq 1$. Then, for any positive integer n, the following expressions*

$$\begin{aligned}
&\sum_{k+l+m=n-1} \binom{n}{k,l,m} \beta^{(l+m+2)}_{k,q^{w_{\sigma(2)}w_{\sigma(3)}}} \left(w_{\sigma(1)}y_1\right) \\
&\times \beta^{(m+2)}_{l,q^{w_{\sigma(1)}w_{\sigma(3)}}} \left(w_{\sigma(2)}y_2\right) T_{2,m}\left(w_{\sigma(3)}-1|q^{w_{\sigma(1)}w_{\sigma(2)}}\right) \\
&\times q^{w_1w_2w_3(l+m+1)y_1} q^{w_1w_2w_3(m+1)y_2} \\
&\times \left[w_{\sigma(2)}w_{\sigma(3)}\right]_q^k \left[w_{\sigma(1)}w_{\sigma(3)}\right]_q^l \left[w_{\sigma(1)}w_{\sigma(2)}\right]_q^{m+1} \\
&+ (q-1) \sum_{k+l+m=n} \binom{n}{k,l,m} \beta^{(l+m+1)}_{k,q^{w_{\sigma(2)}w_{\sigma(3)}}} \left(w_{\sigma(1)}y_1\right) \\
&\times \beta^{(m+1)}_{l,q^{w_{\sigma(1)}w_{\sigma(3)}}} \left(w_{\sigma(2)}y_2\right) T_{1,m}\left(w_{\sigma(3)}-1|q^{w_{\sigma(1)}w_{\sigma(2)}}\right) \\
&\times q^{w_1w_2w_3(l+m)y_1} q^{w_1w_2w_3my_2} \left[w_{\sigma(2)}w_{\sigma(3)}\right]_q^k \left[w_{\sigma(1)}w_{\sigma(3)}\right]_q^l \left[w_{\sigma(1)}w_{\sigma(2)}\right]_q^{m+1}
\end{aligned}$$

are the same for any $\sigma \in S_3$.

Remark. We can get interesting identities by letting $w_3 = 1$ or by letting $w_2 = w_3 = 1$, in view of (2.15). However, writing those down requires much space. So we omit it.

With the same $a = q^{w_2w_3(x_1+w_1y_1)}$, $b = q^{w_1w_3(x_2+w_2y_2)}$ as in (2.18), I_1 can be rewritten as

$$\begin{aligned}
I_1 =& \int_{\mathbb{Z}_p^2} e^{[w_2w_3]_q[x_1+w_1y_1]_{q^{w_2w_3}}t} e^{[w_1w_3]_q[x_2+w_2y_2]_{q^{w_1w_3}}(at)} \\
&\times \left\{ \int_{\mathbb{Z}_p} \left(q^{w_1w_2w_3} e^{[w_1w_2(x_3+w_3)]_q(abt)} - e^{[w_1w_2x_3]_q(abt)} \right) d\mu_{q^{w_1w_2}}(x_3) \right\} \\
&\times d\mu_{q^{w_2w_3}}(x_1)\, d\mu_{q^{w_1w_3}}(x_2).
\end{aligned} \tag{2.25}$$

From Lemma 2 and (2.25), we note that the inner integral is equal to

$$\begin{aligned}
&abt\,[w_1w_2]_q \sum_{i=0}^{w_3-1} q^{2w_1w_2i} e^{[w_1w_2i]_q abt} \\
&+ (q-1)\,[w_1w_2]_q \sum_{i=0}^{w_3-1} q^{w_1w_2i} e^{[w_1w_2i]_q abt}.
\end{aligned} \tag{2.26}$$

Thus, by (2.25) and (2.26), we get

$$\begin{aligned}
I_1 =& t\,[w_1w_2]_q \sum_{i=0}^{w_3-1} q^{2w_1w_2i} \int_{\mathbb{Z}_p^2} ab e^{[w_2w_3]_q[x_1+w_1y_1]_{q^{w_2w_3}}t} \\
&\times e^{[w_1w_3]_q\left[x_2+w_2y_2+\frac{w_2}{w_3}i\right]_{q^{w_1w_3}}at} d\mu_{q^{w_2w_3}}(x_1)\, d\mu_{q^{w_1w_3}}(x_2) \\
&+ (q-1)\,[w_1w_2]_q \sum_{i=0}^{w_3-1} q^{w_1w_2i} \int_{\mathbb{Z}_p^2} e^{[w_2w_3]_q[x_1+w_1y_1]_{q^{w_2w_3}}t} \\
&\times e^{[w_1w_3]_q\left[x_2+w_2y_2+\frac{w_2}{w_3}i\right]_{q^{w_1w_3}}at} d\mu_{q^{w_2w_3}}(x_1)\, d\mu_{q^{w_1w_3}}(x_2)
\end{aligned} \tag{2.27}$$

$$
\begin{aligned}
=&t\left[w_1w_2\right]_q\sum_{i=0}^{w_3-1}q^{2w_1w_2i}\sum_{k,l=0}^{\infty}\frac{t^{k+l}}{k!l!}q^{w_1w_2w_3(l+1)y_1}q^{w_1w_2w_3y_2}\left[w_2w_3\right]_q^k\\
&\times\left[w_1w_3\right]_q^l\int_{\mathbb{Z}_p}q^{w_2w_3(l+1)x_1}\left[x_1+w_1y_1\right]_{q^{w_2w_3}}^k d\mu_{q^{w_2w_3}}\left(x_1\right)\\
&\times\int_{\mathbb{Z}_p}q^{w_1w_3x_2}\left[x_2+w_2y_2+\frac{w_2}{w_3}i\right]_{q^{w_1w_3}}^l d\mu_{q^{w_1w_3}}\left(x_2\right)\\
&+\left(q-1\right)\left[w_1w_2\right]_q\sum_{i=0}^{w_3-1}q^{w_1w_2i}\sum_{k,l=0}^{\infty}\frac{t^{k+l}}{k!l!}q^{w_1w_2w_3ly_1}\left[w_2w_3\right]_q^k\\
&\times\left[w_1w_3\right]_q^l\int_{\mathbb{Z}_p}q^{w_2w_3lx_1}\left[x_1+w_1y_1\right]_{q^{w_2w_3}}^k d\mu_{q^{w_2w_3}}\left(x_1\right)\\
&\times\int_{\mathbb{Z}_p}\left[x_2+w_2y_2+\frac{w_2}{w_3}i\right]_{q^{w_1w_3}}^l d\mu_{q^{w_1w_3}}\left(x_2\right)
\end{aligned}
$$

$$
\begin{aligned}
=&\sum_{n=0}^{\infty}\left\{\sum_{k=0}^{n}\binom{n}{k}\beta_{k,q^{w_2w_3}}^{(n-k+2)}\left(w_1y_1\right)q^{w_1w_2w_3(n-k+1)y_1}\right.\\
&\times q^{w_1w_2w_3y_2}\left[w_1w_2\right]_q\left[w_2w_3\right]_q^k\\
&\left.\times\left[w_1w_3\right]_q^{n-k}\sum_{i=0}^{w_3-1}q^{2w_1w_2i}\beta_{n-k,q^{w_1w_3}}^{(2)}\left(w_2y_2+\frac{w_2}{w_3}i\right)\right\}\frac{t^{n+1}}{n!}\\
&+\left(q-1\right)\sum_{n=0}^{\infty}\left\{\sum_{k=0}^{n}\binom{n}{k}\beta_{k,q^{w_2w_3}}^{(n-k+1)}\left(w_1y_1\right)q^{w_1w_2w_3(n-k)y_1}\right.\\
&\times\left[w_1w_2\right]_q\left[w_2w_3\right]_q^k\left[w_1w_3\right]_q^{n-k}\\
&\left.\times\sum_{i=0}^{w_3-1}q^{w_1w_2i}\beta_{n-k,q^{w_1w_3}}\left(w_2y_2+\frac{w_2}{w_3}i\right)\right\}\frac{t^n}{n!}.
\end{aligned}
$$

Separating the term corresponding to $n=0$ from the second sum and after rearranging, we obtain

$$
\begin{aligned}
I_1=&\left(q-1\right)\left[w_1w_2w_3\right]_q+\sum_{n=1}^{\infty}\left\{n\sum_{k=0}^{n-1}\binom{n-1}{k}\beta_{k,q^{w_2w_3}}^{(n-k+1)}\left(w_1y_1\right)q^{w_1w_2w_3(n-k)y_1}\right.\\
&\times q^{w_1w_2w_3y_2}\left[w_1w_2\right]_q\left[w_2w_3\right]_q^k\left[w_1w_3\right]_q^{n-1-k}\sum_{i=0}^{w_3-1}q^{2w_1w_2i}\\
&\times\beta_{n-1-k,q^{w_1w_3}}^{(2)}\left(w_2y_2+\frac{w_2}{w_3}i\right)+\left(q-1\right)\sum_{k=0}^{n}\binom{n}{k}\beta_{k,q^{w_2w_3}}^{(n-k+1)}\left(w_1y_1\right)q^{w_1w_2w_3(n-k)y_1}\\
&\left.\times\left[w_1w_2\right]_q\left[w_2w_3\right]_q^k\left[w_1w_3\right]_q^{n-k}\sum_{i=0}^{w_3-1}q^{w_1w_2i}\beta_{n-k,q^{w_1w_3}}\left(w_2y_2+\frac{w_2}{w_3}i\right)\right\}\frac{t^n}{n!}.
\end{aligned}
$$

As this expression is invariant under any permutations in w_1, w_2, w_3, we have the following results.

Theorem 4. *Let $w_1, w_2, w_3 \in \mathbb{Z}$ with $w_1 \geq 1$, $w_2 \geq 1$, $w_3 \geq 1$. Then, for any positive integer n, the following expressions*

$$\begin{aligned}
& n\sum_{k=0}^{n-1}\binom{n-1}{k}\beta^{(n-k+1)}_{k,q^{w_{\sigma(2)}w_{\sigma(3)}}}\left(w_{\sigma(1)}y_1\right)q^{w_1w_2w_3(n-k)y_1} \\
& \times q^{w_1w_2w_3y_2}\left[w_{\sigma(1)}w_{\sigma(2)}\right]_q\left[w_{\sigma(2)}w_{\sigma(3)}\right]_q^k\left[w_{\sigma(1)}w_{\sigma(3)}\right]_q^{n-1-k} \\
& \times \sum_{i=0}^{w_{\sigma(3)}-1} q^{2w_{\sigma(1)}w_{\sigma(2)}i}\beta^{(2)}_{n-1-k,q^{w_{\sigma(1)}w_{\sigma(3)}}}\left(w_{\sigma(2)}y_2+\frac{w_{\sigma(2)}}{w_{\sigma(3)}}i\right) \\
& +(q-1)\sum_{k=0}^{n}\binom{n}{k}\beta^{(n-k+1)}_{k,q^{w_{\sigma(2)}w_{\sigma(3)}}}\left(w_{\sigma(1)}y_1\right)q^{w_1w_2w_3(n-k)y_1} \\
& \times\left[w_{\sigma(1)}w_{\sigma(2)}\right]_q\left[w_{\sigma(2)}w_{\sigma(3)}\right]_q^k\left[w_{\sigma(1)}w_{\sigma(3)}\right]_q^{n-k} \\
& \times \sum_{i=0}^{w_{\sigma(3)}-1} q^{w_{\sigma(1)}w_{\sigma(2)}i}\beta_{n-k,q^{w_{\sigma(1)}w_{\sigma(3)}}}\left(w_{\sigma(2)}y_2+\frac{w_{\sigma(2)}}{w_{\sigma(3)}}i\right)
\end{aligned}$$

are all the same for any $\sigma \in S_3$.

Remark. Using (2.14) and by specializing $w_3 = 1$ or $w_2 = w_3 = 1$, we can obtain many interesting identities. However, as this requires much space, we will omit those.

ACKNOWLEDGEMENT

This work was supported by the National Research Foundation of Korea(NRF) grant funded by the Korea government(MOE) (No.2012R1A1A2003786) and partially supported by Kwangwoon University 2014.

REFERENCES

- [1] M. Açikgöz, D. Erdal, and S. Araci, *A new approach to q-Bernoulli numbers and q-Bernoulli polynomials related to q-Bernstein polynomials*, Adv. Difference Equ. (2010), Art. ID 951764, 9. MR 2764538 (2012g:11038)
- [2] R. P. Agarwal and C. S. Ryoo, *Numerical computations of the roots of the generalized twisted q-Bernoulli polynomials*, Neural Parallel Sci. Comput. **15** (2007), no. 2, 193–206. MR 2357405 (2008i:65095)
- [3] A. Bayad, *Special values of Lerch zeta function and their Fourier expansions*, Adv. Stud. Contemp. Math. (Kyungshang) **21** (2011), no. 1, 1–4. MR 2984857
- [4] L. Carlitz, *q-Bernoulli numbers and polynomials*, Duke Math. J. **15** (1948), 987–1000. MR 0027288 (10,283g)
- [5] ______, *Expansions of q-Bernoulli numbers*, Duke Math. J. **25** (1958), 355–364. MR 0095982 (20 #2480)
- [6] J. Choi, T. Kim, and Y. H. Kim, *A note on the extended q-Bernoulli numbers and polynomials*, Adv. Stud. Contemp. Math. (Kyungshang) **21** (2011), no. 4, 351–354. MR 2884998 (2012j:11045)
- [7] D. Ding and J. Yang, *Some identities related to the Apostol-Euler and Apostol-Bernoulli polynomials*, Adv. Stud. Contemp. Math. (Kyungshang) **20** (2010), no. 1, 7–21. MR 2597988 (2011k:05030)

[8] D. V. Dolgy and T. Kim, *A note on the weighted q-Bernoulli numbers and the weighted q-Bernstein polynomials*, Honam Math. J. **33** (2011), no. 4, 519–527. MR 2906484 (2012k:11021)

[9] Y. He, *Symmetric identities for Carlitz's q-Bernoulli numbers and polynomials*, Adv. Difference Equ. (2013), 2013:246, 10. MR 3116122

[10] K.-W. Hwang, D. V. Dolgy, D. S. Kim, T. Kim, and S. H. Lee, *Some theorems on Bernoulli and Euler numbers*, Ars Combin. **109** (2013), 285–297. MR 3087218

[11] D. S. Kim, N. Lee, J. Na, and K. H. Park, *Abundant symmetry for higher-order Bernoulli polynomials (I)*, Adv. Stud. Contemp. Math. (Kyungshang) **23** (2013), no. 3, 461–482. MR 3113161

[12] G. Kim, B. Kim, and J. Choi, *The DC algorithm for computing sums of powers of consecutive integers and Bernoulli numbers*, Adv. Stud. Contemp. Math. (Kyungshang) **17** (2008), no. 2, 137–145. MR 2452155

[13] T. Kim, *q-Volkenborn integration*, Russ. J. Math. Phys. **9** (2002), no. 3, 288–299. MR 1965383 (2004f:11138)

[14] ______, *Sums of powers of consecutive q-integers*, Adv. Stud. Contemp. Math. (Kyungshang) **9** (2004), no. 1, 15–18. MR 2067834 (2005c:11152)

[15] ______, *Power series and asymptotic series associated with the q-analog of the two-variable p-adic L-function*, Russ. J. Math. Phys. **12** (2005), no. 2, 186–196. MR 2198999 (2007d:11102)

[16] ______, *q-Bernoulli numbers and polynomials associated with Gaussian binomial coefficients*, Russ. J. Math. Phys. **15** (2008), no. 1, 51–57. MR 2390694 (2009b:11040)

[17] ______, *Symmetry p-adic invariant integral on $\mathbb{Z}_p$ for Bernoulli and Euler polynomials*, J. Difference Equ. Appl. **14** (2008), no. 12, 1267–1277. MR 2462529 (2009i:11023)

[18] ______, *Symmetry of power sum polynomials and multivariate fermionic p-adic invariant integral on $\mathbb{Z}_p$*, Russ. J. Math. Phys. **16** (2009), no. 1, 93–96. MR 2486809 (2010c:11028)

[19] ______, *Some identities for the Bernoulli, the Euler and the Genocchi numbers and polynomials*, Adv. Stud. Contemp. Math. (Kyungshang) **20** (2010), no. 1, 23–28. MR 2597989 (2011a:11211)

[20] ______, *A family of (h,q)-zeta function associated with (h,q)-Bernoulli numbers and polynomials*, J. Comput. Anal. Appl. **14** (2012), no. 3, 402–409. MR 2932567

[21] T. Kim, C. S. Ryoo, and H. Yi, *A note on q-Bernoulli numbers and q-Bernstein polynomials*, Ars Combin. **104** (2012), 437–447. MR 2951805

[22] Y.-H. Kim and K.-W. Hwang, *Symmetry of power sum and twisted Bernoulli polynomials*, Adv. Stud. Contemp. Math. (Kyungshang) **18** (2009), no. 2, 127–133. MR 2508978 (2010b:11027)

[23] Q.-M. Luo, *Some recursion formulae and relations for Bernoulli numbers and Euler numbers of higher order*, Adv. Stud. Contemp. Math. (Kyungshang) **10** (2005), no. 1, 63–70. MR 2108686 (2005h:11039)

[24] H. Ozden, I. N. Cangul, and Y. Simsek, *Remarks on q-Bernoulli numbers associated with Daehee numbers*, Adv. Stud. Contemp. Math. (Kyungshang) **18** (2009), no. 1, 41–48. MR 2479746 (2009k:11037)

[25] S.-H. Rim, A. Bayad, E.-J. Moon, J.-H. Jin, and S.-J. Lee, *A new construction on the q-Bernoulli polynomials*, Adv. Difference Equ. (2011), 2011:34, 6. MR 2844137 (2012h:11025)

[26] S.-H. Rim and T. Kim, *Explicit p-adic expansion for alternating sums of powers*, Adv. Stud. Contemp. Math. (Kyungshang) **14** (2007), no. 2, 241–250. MR 2316990 (2008k:11126)

[27] S.-H. Rim, T. Kim, and B.-J. Lee, *Some identities on the extended Carlitz's q-Bernoulli numbers and polynomials*, J. Comput. Anal. Appl. **14** (2012), no. 3, 536–543. MR 2933541

[28] C. S. Ryoo, T. Kim, and B. Lee, *q-Bernoulli numbers and q-Bernoulli polynomials revisited*, Adv. Difference Equ. (2011), 2011:33, 8. MR 2844136 (2012h:11026)

[29] Y. Simsek, *Theorems on twisted L-function and twisted Bernoulli numbers*, Adv. Stud. Contemp. Math. (Kyungshang) **11** (2005), no. 2, 205–218. MR 2169895 (2006e:11123)

[30] ———, *Generating functions of the twisted Bernoulli numbers and polynomials associated with their interpolation functions*, Adv. Stud. Contemp. Math. (Kyungshang) **16** (2008), no. 2, 251–278. MR 2404639 (2009f:11021)

[31] H. M. Srivastava, T. Kim, and Y. Simsek, *q-Bernoulli numbers and polynomials associated with multiple q-zeta functions and basic L-series*, Russ. J. Math. Phys. **12** (2005), no. 2, 241–268. MR 2199005 (2006k:11034)

Institute of Mathematics and Computer Sciences, Far Eastern Federal University, Vladivostok 690060, Russia
E-mail address: *d_dol@mail.ru*

Department of Mathematics, Sogang University, Seoul 121-742, Republic of Korea
E-mail address: `dskim@sogang.ac.kr`

Department of Mathematics, Kwangwoon University, Seoul 139-701, Republic of Korea
E-mail address: `tkkim@kw.ac.kr`